

Generalized Tsen theorem and rationally connected Fano fibrations

Frederic Campana, Thomas Peternell and Aleksandr Pukhlikov

We prove that a fibration $X/\mathbb{P}^1$ the general fiber of which is a smooth Fano three-fold is rationally connected. The proof is based upon a generalization of the classical Tsen theorem: we prove that a fibration X/C over a curve C the general fiber of which is a Fano complete intersection in a product of (weighted) projective spaces has a section.

1 Introduction

This paper is concerned with the following problem

Problem 1.1 *Let $\pi: X \rightarrow \mathbb{P}^1$ be a fibration, the general fiber of which is rationally connected. Then X is rationally connected.*

A positive solution of this problem has the important consequence that fibrations over rationally connected varieties with rationally connected fiber are again rationally connected. If one accepts the predictions of Mori theory, then one is reduced to the analysis of Fano fibrations over $\mathbb{P}^1$, however one should have in mind that the fibers may all be singular in this approach since the total space is singular in general when passing from the original X to a model admitting a Fano fibration.

We have two main results:

1.1 Generalized Tsen theorem

Let $\mathbb{P}$ denote the weighted projective space $\mathbb{P}(a_0, a_1, \dots, a_m)$. Let $\mathcal{V}$ denote the class of Fano complete intersections in the product of weighted projective spaces $\mathbb{P}^{(1)} \times \dots \times \mathbb{P}^{(k)} = \mathbb{P}^\times$ where $\mathbb{P}^{(i)} = \mathbb{P}(a_0^{(i)}, a_1^{(i)}, \dots, a_{m(i)}^{(i)})$

Theorem 1.2 (the generalized Tsen theorem) *Let $\pi: X \rightarrow C$ be a fibration over a curve, the general fiber of which belongs to the class $\mathcal{V}$. Then there exists a section $\beta: C \rightarrow X$, $\pi \circ \beta = \text{id}_C$.*

Corollary 1.3 *Let $\pi: X \rightarrow \mathbb{P}^1$ be a fibration over a rational curve, the general fiber of which is a Fano complete intersection in the product of weighted projective spaces. Then X is rationally connected.*

1.2 Fano fibrations

Theorem 1.4 *Let $\pi: X \rightarrow \mathbb{P}^1$ be a fibration, the general fiber of which is a smooth Fano threefold. Then X is rationally connected.*

The proof of this theorem is by running through the classification of Fano 3—folds by Iskovskikh, Mori and Mukai and using ad hoc methods in the single cases. We are giving details only in case $b_2 = 1$ while details for $b_2 \geq 2$ will be written up at a different place.

After having completed our work up to some redigations, we got aware of the new preprint [GHS01] in which the problem 1.1 appears to be completely solved with totally different methods from deformation theory and moduli theory of curves in the meantime also a preprint of de Jong-Starr appeared. However this paper remains meaningful, because the methods presented here are effective (sections are explicitly constructed) and they seem to be much simpler. An interesting aspect is also that our approach avoids the involved considerations of multiple components of fibers.

1.3 Plan of the proof of the generalized Tsen theorem

We will prove a particular case of Theorem 2, when $\mathbb{P}^{(i)} \equiv \mathbb{P} = \mathbb{P}^M$ is the ordinary projective space and the general fiber $V \subset \mathbb{P} \times \dots \times \mathbb{P}$ is a hypersurface of the type $(M, \dots, M)$. In the general case no modification of the method is necessary, all the technique developed below works automatically as it is (up to the Lefschetz theorem 1.5). However, in the general case one needs considerably more complicated notations. This is essentially the only reason why we restrict ourselves by this typical particular case.

Our proof consists of two main steps:

- first, we construct a special model $X^\sharp/C$, birational to X/C over C :

$$\begin{array}{ccc} X & \xrightarrow{\chi} & X^\sharp \\ \downarrow & & \downarrow \\ C & = & C, \end{array}$$

where over a point $x \in C$ of general position the birational map χ is a *biregular* isomorphism of fibers, so that, in particular, χ induces a bijection

$$\text{Sect}(X/C) \leftrightarrow \text{Sect}(X^\sharp/C)$$

of the sets of sections and therefore the problems of existence of a section of X/C and $X^\sharp/C$ are equivalent.

Let us explain the meaning of this step. By assumption, a fiber of general position of the fibration X/C permits an embedding

$$X_x = \pi^{-1}(x) \subset \mathbb{P} \times \dots \times \mathbb{P}$$

as hypersurface of type $(M, \dots, M)$ in $\mathbb{P} \times \dots \times \mathbb{P} = \mathbb{P}^{\times k}$. However, in the general case this embedding cannot be globalized, even on the birational level, that is, there exists no birational map over C

$$X \dashrightarrow \mathbb{P}^{\times k} \times C$$

to the trivial (or a locally trivial) $\mathbb{P}^{\times k}$ -fibration over C , which would have embedded a fiber of general position into $\mathbb{P}^{\times k}$ as a hypersurface of the type $(M, \dots, M)$. It is exactly for this reason that the classical Tsen theorem [Ts36, Ko96] does not work. We will show, however (Sec. 2), that there exists a certain analogue of such a globalization: an embedding

$$\begin{array}{ccc} X^\sharp & \hookrightarrow & \mathbb{P}^\sharp \\ & & \downarrow \\ & & C \end{array}$$

over C , where $\mathbb{P}^\sharp$ is fibered over C with the general fiber $\mathbb{P}^{\times k}$ (but of course $\mathbb{P}^\sharp/C$ is not a locally trivial $\mathbb{P}^{\times k}$ -fibration). The construction of the space $\mathbb{P}^\sharp$ and the model $X^\sharp$ is very explicit and makes it possible to study the problem of existence of sections.

- Second, we prove existence of sections of $X^\sharp/C \hookrightarrow \mathbb{P}^\sharp/C$ (and simultaneously give a description of all sections). We proceed in two steps:
 - to begin with, we describe the sections of the ambient fibration $\mathbb{P}^\sharp/C$.
 - After that, using the explicit description of those divisors $X^\sharp \subset \mathbb{P}^\sharp$, the fiber of which over a point of general position $x \in C$ is a $(M, \dots, M)$ -hypersurface in $\mathbb{P}^{\times k}$, we make the crucial step: we write out the conditions (a system of algebraic equations on the parameters, defining the sections of the fibration $\mathbb{P}^\sharp/C$), which make a section $s: C \rightarrow \mathbb{P}^\sharp$ lie where it should:

$$s(C) \subset X^\sharp.$$

Now a simple dimension count in the spirit of the classical Tsen theorem [Ts36], [Ko96] shows that these conditions really can be satisfied, which completes the proof.

There is one point to be a little careful concerning complete intersections in weighted projective spaces:

Proposition 1.5 *Let $\mathbb{P}^j$ be weighted projective spaces, $\mathbb{P} = \mathbb{P}^1 \times \dots \times \mathbb{P}^r$ and $X \subset \mathbb{P}$ be a quasi-smooth complete intersection of dimension at least 3. Then the restriction map*

$$\mathrm{Pic}(\mathbb{P}) \rightarrow \mathrm{Pic}(X)$$

is an isomorphism.

Proof. Of course this is the classical Lefschetz theorem in case $\mathbb{P}$ and X are smooth. In general there are three ways to conclude.

- (a) Adapt the method of [Do81,3.2.4/3.2.5] (where the case of one weighted projective space is treated).
- (b) Use [Do80] (see remark (3.2.6) in [Do81]).
- (c) Argue directly as follows. First we consider general quasi-smooth complete intersections X . Let $\mathbb{P}_{m_j} \rightarrow \mathbb{P}^j$ the quotient map, let $\tilde{\mathbb{P}} = \mathbb{P}_{m_1} \times \dots \times \mathbb{P}_{m_r}$ and consider the preimage $\tilde{X}$ of X in $\tilde{\mathbb{P}}$. Since X is general, $\tilde{X}$ will be smooth. Now apply the Lefschetz theorem for $\tilde{X} \subset \tilde{\mathbb{P}}$ and go back to X . Of course this works only for general X . But any quasi-smooth X is a deformation of those X for which we already know our claim. Then we see immediately that the claim holds also for the special X : it is of course clear that $\text{Pic}(X) = \text{Pic}(X_t)$, and a small argument shows that every line bundle on X is induced by a line bundle on $\mathbb{P}$, since it is topologically the case and the analytic cohomology in the exponential sequence vanishes. Q.E.D.

1.4 Acknowledgements

For the third author, the present research has been carried out in the framework of his Humboldt research project, and he is very thankful to the Humboldt Foundation for the financial support. The first author was supported by the Schwerpunktprogramm “Global methods in complex geometry” of the Deutsche Forschungsgemeinschaft.

2 Construction of the special model

In this section we realize the first step of our program: we construct a special model of the fibration X/C , adjusted to our program.

2.1 The finite cover, parametrizing projections of the fibers

Consider the fibration X/C as in subsection 1.3. For a point $x \in C$ of general position the fiber $\pi^{-1}(x) = X_x$ is a hypersurface of the type $(M, \dots, M)$ in $\mathbb{P}^{\times k}$. Therefore, for X_x one gets k canonically given projections

$$\text{pr}: X_x \rightarrow \mathbb{P}.$$

These k projections can be reconstructed from the variety X_x itself and do not depend on its particular projective embedding.

Proposition 2.1 *There exists a projective curve B (reduced, but possibly reducible), which is a k -sheeted cover of the curve C :*

$$\tau: B \rightarrow C,$$

$\deg \tau = k$, the points of which parametrize the projections of the fibers $X_x \rightarrow \mathbb{P}$ in the following sense. There exists a regular algebraic map

$$p: \pi^{-1}(C^\circ) \times_C B \rightarrow \mathbb{P}, \quad (1)$$

where $C^\circ \subset C$ is an open set of all such points $x \in C$ that the fiber X_x is a hypersurface of type $(M, \dots, M)$ in $\mathbb{P}^{\times k}$ (recall that by assumption this is true for a fiber of general position only), the fiber $\tau^{-1}(x) \subset B$ consists of precisely k distinct points,

$$\tau^{-1}(x) = \{y_1, \dots, y_k\},$$

and the k morphisms

$$p|_{X_x \times \{y_i\}}: X_x \times \{y_i\} \rightarrow \mathbb{P}$$

are precisely k projections

$$\text{pr}: X_x \rightarrow \mathbb{P} \times \dots \times \mathbb{P} \rightarrow \mathbb{P}$$

onto the factors for each point $x \in C^\circ$.

Proof. For simplicity let $k = 2$. The fibers of the projections $X_x \rightarrow \mathbb{P}^i$, $x \in C$, $i = 1, 2$ define either two or one irreducible families of cycles. In the first case we obtain two rational maps $X \rightarrow \mathbb{P}^i$, which are regular on the general X_x . Then we let B be two disjoint copies of C and the rest is clear.

So we may suppose that we have one irreducible family; let $\mathcal{C} \rightarrow T$ denote its graph with projection $\rho: \mathcal{C} \rightarrow X$. Then we obtain canonically a map $h: T \rightarrow C$ which is of course a 2-sheeted cover (generically). Therefore we set $B = T$. Over C^0 , we have $\mathcal{C} = X \times_{C^0} B^0$, and there exists a map $\mathcal{C} \rightarrow \mathbb{P}$, just by sending $z \in C_t$ to the point in P , to which $\rho(C_t)$ is mapped by the corresponding projection $X_h(z) \rightarrow \mathbb{P}^i$. Q.E.D.

2.2 Construction of the ambient space $\mathbb{P}^\sharp = \mathbb{P}^{B/C}$: an analytic description over C°

Denote by the symbol $\mathbb{P}_B = \mathbb{P} \times B$ the trivial $\mathbb{P}$ -fibration over B , and by $\pi_B: \mathbb{P}_B \rightarrow B$ the corresponding projection. Let $\mathbb{P}_y = \mathbb{P} \times \{y\} = \pi_B^{-1}(y)$ be the fiber over a point $y \in B$, and $\pi_C: \mathbb{P}_B \rightarrow C$ be the composition $\pi_C = \tau \circ \pi_B$. As above, denote by $C^\circ \subset C$, the set of points of general position, which have precisely k preimages in B .

An open (in the real topology) set $D \subset C^\circ$ (for instance, a small disk) is said to be *simple*, if

$$\tau^{-1}(D) = \coprod_{i=1}^k D_i,$$

where $\tau: D_i \rightarrow D$ is an isomorphism for each i . Now set for a simple $D \subset C^\circ$:

$$\mathbb{P}^{B/C}(D) = \prod_{i=1}^k \mathbb{P}_D(\mathbb{P} \times D_i) \cong \mathbb{P}^{\times k} \times D,$$

where the symbol $\prod_D$ stands for the fiber product with respect to the morphisms

$$\mathbb{P} \times D_i \rightarrow D_i \rightarrow D.$$

Let

$$\kappa_i: \mathbb{P}^{B/C}(D) \rightarrow \mathbb{P}$$

mean the projection onto the i -th factor. Now set $X(D) = \pi^{-1}(D) \subset X$ and look at the fiber product

$$X(D) \times_C B = X(D) \times_C \tau^{-1}(D) \simeq \prod_{i=1}^k X(D)^{(i)},$$

where $X(D)^{(i)}$ is the identical copy of the set $X(D)$, corresponding to the connected component D_i . Let

$$\nu_i: X(D) \rightarrow X(D)^{(i)}$$

be the corresponding isomorphism.

Lemma 2.2 *There exists a uniquely determined natural morphism — an inclusion*

$$\iota_D: X(D) \rightarrow \mathbb{P}^{B/C}(D)$$

such that for each $i = 1, \dots, k$ the following diagram is commutative:

$$\begin{array}{ccccc} X(D) & \xrightarrow{\iota_D} & \mathbb{P}^{B/C}(D) & & \\ \nu_i \downarrow & & \downarrow & \kappa_i & \\ X(D)^{(i)} & \xrightarrow{p_D} & \mathbb{P}, & & \end{array} \quad (2)$$

where

$$p_D: X(D) \times_C B \rightarrow \mathbb{P} \quad (3)$$

is given by $p|_D \circ pr_D$, with $p|_D$ the restriction of the projection p from Proposition 2.1 onto the set D .

Proof. Define the morphism ι_D by the formula

$$\iota_D = \prod_{i=1}^k (p \circ \nu_i) \times \pi: X(D) \rightarrow \mathbb{P}^{B/C}(D) = \left(\prod_{i=1}^k \mathbb{P} \right) \times D.$$

The diagram (2) is now obviously commutative. It is also obvious that (3) determines the morphism ι_D uniquely. For a fiber X_x , $x \in D$, it means simply that the k projections

$$p_y: X_x \rightarrow \mathbb{P}_y,$$

$y \in \tau^{-1}(x)$, determine the embedding

$$\iota_x: X_x \rightarrow \prod_{y \in \tau^{-1}(x)} \mathbb{P}_y.$$

Q.E.D.

Remark 2.3 Let $D^\sharp \subset D$ be an open subset. Then $D^\sharp$ is simple, too:

$$\tau^{-1}(D^\sharp) = \coprod_{i=1}^k D_i^\sharp,$$

so that the construction, described above, can be applied to $D^\sharp$, too. It is easy to see that there is a uniquely determined natural inclusion

$$\iota_{D^\sharp, D}: \mathbb{P}^{B/C}(D^\sharp) \hookrightarrow \mathbb{P}^{B/C}(D),$$

which makes the following diagram commutative:

$$\begin{array}{ccc} X(D^\sharp) & \hookrightarrow & X(D) \\ \downarrow \iota_{D^\sharp} & & \downarrow \iota_D \\ \mathbb{P}^{B/C}(D^\sharp) & \hookrightarrow & \mathbb{P}^{B/C}(D). \end{array}$$

This makes it possible to glue the varieties $\mathbb{P}^{B/C}(D)$ and the inclusions $\iota_D: X(D) \hookrightarrow \mathbb{P}^{B/C}(D)$ together and obtain an (open) variety $\mathbb{P}^{B/C}(C^\circ)$ and an embedding $X(C^\circ) \subset \mathbb{P}^{B/C}(C^\circ)$.

2.3 Construction of the ambient space $\mathbb{P}^{B/C}$: an analytic description in the ramification case

To be able to work conveniently with the space $\mathbb{P}^{B/C}$ it is necessary to complete it over the points $x \in C \setminus C^\circ$. We may assume that if the curve B has several irreducible components, then they are disjoint, that is, if a fiber $\tau^{-1}(x)$ consists of less than k points, then it happens due to ramification of the morphism τ . We also may assume that the exceptional set $C \setminus C^\circ$ consists of such points only, because if $\sharp\tau^{-1}(x) = k$, then the space $\mathbb{P}^{B/C}$ is constructed exactly as above (although the embedding morphism i for the corresponding fiber $X_x \hookrightarrow \mathbb{P}_x^{B/C}$ is possibly no longer defined).

Now take a point $x \in C \setminus C^\circ$ and consider

$$\tau^*x = \sum_{i=1}^a l_i y_i,$$

where $a \leq k-1$, $l_i \geq 1$ and at least one of the integers l_i is strictly bigger than 1. Let $D \subset C$ be a small disk with the centre at the point x , where

$$\tau^{-1}(D) = \coprod_{i=1}^a D_i, \quad D_i \ni y_i,$$

and there exist local coordinates z on D and z_i on D_i , $z(x) = z_i(y_i) = 0$, respectively, such that the morphism

$$\tau: D_i \rightarrow D$$

takes the form

$$\tau: z_i \mapsto z_i^{l_i}.$$

Let ζ_i be a primitive root of degree l_i from 1 and $G_i = \mathbb{Z}_{l_i}$ be the cyclic group of order l_i .

Set

$$\begin{aligned} Z(D) &= \prod_{i=1}^a \left(\mathbb{P}^{\times l_i} \times D_i \right) = \\ &= \{ ((p_1^{(1)}; \dots; p_{l_1}^{(1)}; z_1), (p_1^{(2)}; \dots; p_{l_2}^{(2)}; z_2), \dots, (p_1^{(a)}; \dots; p_{l_a}^{(a)}; z_a)) \mid z_1^{l_1} = \dots = z_a^{l_a} \}. \end{aligned}$$

Now we define $\mathbb{P}^{B/C}$ locally on D in the following way:

$$\mathbb{P}^{B/C}(D) = Z(D)/G_1 \times \dots \times G_a,$$

where the cyclic group $G_i = \langle \sigma_i \rangle$ acts on the i -th factor of the fiber product $Z(D)$ as follows

$$\sigma_i(p_1^{(i)}; \dots; p_{l_i}^{(i)}; z_i) = (p_{l_i}^{(i)}; p_1^{(i)}; \dots; p_{l_i-1}^{(i)}; \zeta_i z_i)$$

(the points $p_j^{(i)}$ are permuted in the cyclic way). This action is obviously compatible with the construction of the fiber product ($(\zeta_i z_i)^{l_i} = z_i^{l_i}$) and thus the defined action of G_i can be naturally extended to an action on $Z^{\text{an}}(D)$ (on the other fiber factors the group G_i acts trivially).

It is checked easily that in the punctured disk $D^\# = D \setminus \{x\}$ this construction coincides with that of Sec. 2.2: more precisely, if $D^\# \subset D$ is a simple open set, so that

$$\tau^{-1}(D^\#) = \coprod_{e=1}^k D_e^\#$$

(in particular, $x \notin D^\#$), then there is a uniquely determined natural inclusion

$$\mathbb{P}^{B/C}(D^\#) \hookrightarrow \mathbb{P}^{B/C}(D),$$

where on the left-hand side we have the space, constructed in Sec. 2.2. Indeed,

$$\tau^{-1}(D^\#) \cap D_i = \prod_{j=1}^{l_i} D_{i,j}^\#,$$

so that over $D^\#$ all the maps

$$\tau_{i,j}: D_{i,j}^\# \rightarrow D^\#$$

are isomorphisms. In particular, one can take l_i single-valued branches of the function $z^{1/l_i} = \tau_{i,j}^{-1}$. Fix one of them and set

$$\mu: D_i \cap \tau^{-1}(D^\#) \rightarrow D_i^\#,$$

$$\mu: z_i \mapsto z_i^\# = (z_i^{l_i})^{1/l_i},$$

where $D_i^\# = D_{i,c(i)}^\#$ is the subset (connected component) $D_{i,j}^\#$, which corresponds to the chosen branch. Now let

$$\lambda: \prod_{i=1}^a D \left(\mathbb{P}^{\times l_i} \times (D_i \cap \tau^{-1}(D^\#)) \right) = Z(D) \cap \left(\prod_{i=1}^a \mathbb{P}^{\times l_i} \cap \tau^{-1}(D^\#) \right) \rightarrow \left(\prod_{i=1}^a \prod_{j=1}^{l_i} \mathbb{P} \right) \times D$$

be given on the i -th factor by the formula

$$(p_1^{(i)}; \dots; p_{l_i}^{(i)}; z_i) \mapsto \sigma_i^f(p_1^{(i)}; \dots; p_{l_i}^{(i)}; z_i),$$

where $f \in \mathbb{Z}$ is defined by the condition

$$z_i^\# = \zeta_i^f z_i.$$

It is easy to check that λ determines an isomorphism

$$\mathbb{P}^{B/C}(D) \cap \pi^{-1}(D^\#) \cong \mathbb{P}^{B/C}(D^\#),$$

which, moreover, does not depend on the choice of ζ_i and the single-valued branch of the root $(\cdot)^{1/l_i}$. This isomorphism identifies $\mathbb{P}^{B/C}(D^\#)$ in the sense of Sec. 2.2 with an open subset of the set $\mathbb{P}^{B/C}(D)$.

What has been just said means that the open pieces $\mathbb{P}^{B/C}(D)$, $D \subset C$, glue together into a compact complex space $\mathbb{P}^{B/C} = \mathbb{P}^{B/C}(C)$ with a natural projection

$$\pi: \mathbb{P}^{B/C} \rightarrow C.$$

Here for any point $x \in C^\circ$ the fiber of the projection π is

$$\mathbb{P}_x^{B/C} = \pi^{-1}(x) = \prod_{y \in \tau^{-1}(x)} \mathbb{P}_y.$$

2.4 Construction of the ambient space $\mathbb{P}^{B/C}$: an algebraic description

The analytic description of $\mathbb{P}^{B/C} = \mathbb{P}_{\text{an}}^{B/C}$ given above is very convenient for local computations. However, a priori it is unclear whether $\mathbb{P}^{B/C}$ is an algebraic variety. So now we give a description of the space $\mathbb{P}^{B/C}$ from the viewpoint of algebraic geometry, which we will denote by $\mathbb{P}^{B/C} = \mathbb{P}_{\text{alg}}^{B/C}$; a little later we will prove that there is an isomorphism of complex spaces

$$\mathbb{P}_{\text{an}}^{B/C} \cong \mathbb{P}_{\text{alg}}^{B/C}.$$

Set

$$\mathbb{P}_B^{\times C^k} = \prod_{i=1}^k {}_C\mathbb{P}_B = \underbrace{\mathbb{P}_B \times_C \mathbb{P}_B \times_C \cdots \times_C \mathbb{P}_B}_k$$

with the natural projection π_B onto

$$B^{\times_C k} = \prod_{i=1}^k {}_C B = \underbrace{B \times_C B \times_C \cdots \times_C B}_k,$$

induced by the projection π_B on each factor.

Let

$$\tau: B^{\times_C k} \rightarrow C^{\times_C k} = \prod_{i=1}^k {}_C C = C$$

be the natural projection. Obviously, $\tau^{-1}(C^\circ)$ is a smooth affine algebraic variety. For an open (in any topology) subset $D \subset C^\circ$ set

$$Z_B(D) = \{(y_1, \dots, y_k) | \tau(y_1) = \dots = \tau(y_k) = x \in D, y_i \neq y_j \text{ for } i \neq j\}.$$

Let

$$Z_B \subset B^{\times_C k}$$

be the closure of the open set $Z_B(C^\circ)$. On the varieties $B^{\times_C k}$ and $\mathbb{P}_B^{\times_C k}$ we have the canonical action of the symmetric group S_k : it permutes the factors of the fiber product. This action is obviously compatible with the projection π_B . For an open set $D \subset C^\circ$ let

$$Z_{\mathbb{P}}(D) = \pi_B^{-1}(Z_B(D)) \subset \mathbb{P}_B^{\times_C k},$$

and let

$$Z_{\mathbb{P}} \subset \mathbb{P}_B^{\times_C k}$$

be the closure of the open set $Z_{\mathbb{P}}(C^\circ)$. The closed sets $Z_B, Z_{\mathbb{P}}$ are invariant with respect to S_k , $\pi_B(Z_{\mathbb{P}}) = Z_B$, $\pi_B^{-1}(Z_B) = Z_{\mathbb{P}}$, and the projection π_B is compatible with the action of the group S_k . It is easy to see that the quotient variety satisfies

$$B^{B/C} = Z_B/S_k \cong C.$$

Now set

$$\mathbb{P}_{\text{alg}}^{B/C} = Z_{\mathbb{P}}/S_k.$$

Proposition 2.4 *The algebraic variety $\mathbb{P}_{\text{alg}}^{B/C}$ and the compact complex space $\mathbb{P}_{\text{an}}^{B/C}$ are canonically isomorphic (as complex spaces).*

Proof. Let $D \subset C^\circ$ be a simple open set, so that over D the cover $\tau: B \rightarrow C$ is reducible,

$$\tau^{-1}(D) = \coprod_{i=1}^k D_i.$$

As above, the symbol π denotes the natural projection for each of the varieties under consideration. By definition,

$$Z_{\mathbb{P}}(D) = \{((p_1, y_1), (p_2, y_2), \dots, (p_k, y_k)) | \tau(y_1) = \dots = \tau(y_k) = x \in D, y_i \neq y_j \text{ for } i \neq j\}.$$

Consider the map

$$\lambda: Z_{\mathbb{P}}(D) \rightarrow \mathbb{P}_{\text{an}}^{B/C}(D)$$

defined in the following way:

$$((p_1, y_1), \dots, (p_k, y_k)) \mapsto (p_{\sigma(1)}; \dots; p_{\sigma(k)}; z = \tau(y_1)) \in \mathbb{P}^{\times k} \times D,$$

where the permutation $\sigma \in S_k$ is uniquely determined by the condition

$$y_{\sigma(i)} \in D_i.$$

It is easy to see that the fibers of the map λ are precisely the orbits of the group S_k , so that λ defines a natural isomorphism

$$Z_{\mathbb{P}}(D)/S_k \cong \mathbb{P}_{\text{an}}^{B/C}(D).$$

Since this isomorphism is natural, it extends to an isomorphism

$$Z_{\mathbb{P}}(C^\circ)/S_k \cong \mathbb{P}_{\text{an}}^{B/C}(C^\circ).$$

We must check that this isomorphism of complex spaces extends to a global isomorphism

$$Z_{\mathbb{P}}/S_k \cong \mathbb{P}_{\text{an}}^{B/C}.$$

To avoid overcomplicated notations, we perform this work for an “extremal” particular case of the maximal ramification, when B has only one branch over $x \in C$. In the general case of several branches the arguments are completely similar, only there are more indices and the formulas are more complicated. So assume that $\tau^{-1}(x) = \{y\}$ and with respect to suitable local coordinates z and w on C and B the map τ is given by the formula

$$\tau^{-1}(D) \ni w \mapsto z = w^k \in D.$$

Recall that

$$\mathbb{P}_{\text{an}}^{B/C}(D) = \{(p_1; \dots; p_k; z)\}/\mathbb{Z}_k,$$

$$\mathbb{P}_{\text{alg}}^{B/C}(D) = Z_{\mathbb{P}}(D)/S_k,$$

where $Z_{\mathbb{P}}(D)$ is the closure of

$$Z_{\mathbb{P}}(D^\circ) = \{((p_1, z_1), \dots, (p_k, z_k)) | z_1^k = \dots = z_k^k = z \in D^\circ, z_i \neq z_j \text{ for } i \neq j\},$$

$D^\circ = D \setminus \{x\}$ being a punctured disk.

Let the symbol $S^\circ(\mathbb{Z}_k)$ denote the set (group) of permutations of the additive group $\mathbb{Z}_k$, which maps 0 to itself. We get the decomposition

$$Z_{\mathbb{P}}(D^\circ) = \coprod_{\sigma \in S^\circ(\mathbb{Z}_k)} Z_{\mathbb{P}}(D^\circ, \sigma),$$

where

$$Z_{\mathbb{P}}(D^{\circ}, \sigma) = \{((p_1, z_1), \dots, (p_k, z_k)) | z_i = \zeta^{\sigma(i-1)} z_1\},$$

ζ is a fixed primitive root of unit of degree k . The group S_k permutes the components $Z_{\mathbb{P}}(D^{\circ}, \sigma)$:

$$\mu: Z_{\mathbb{P}}(D^{\circ}, \sigma) \rightarrow Z_{\mathbb{P}}(D^{\circ}, \mu[\sigma]),$$

$\mu \in S_k$, where the permutation $\mu[\sigma] \in S^{\circ}(\mathbb{Z}_k)$ is defined by the relation

$$\mu[\sigma](i-1) = \sigma(\mu(i)-1) - \sigma(\mu(1)-1), \quad (4)$$

$i = 1, \dots, k$, and we identify the integer i and its image in $\mathbb{Z}_k$. Now set $Z_{\mathbb{P}}(D, \sigma)$ to be the closure of $Z_{\mathbb{P}}(D^{\circ}, \sigma)$ in $Z_{\mathbb{P}}(D)$. Obviously, the group S_k permutes isomorphically the subsets $Z_{\mathbb{P}}(D, \sigma)$ and acts transitively on the set of these subsets. Therefore, if $N(\sigma) \subset S_k$ is the stabilizing subgroup of the component $Z_{\mathbb{P}}(D, \sigma)$, then

$$\mathbb{P}_{\text{alg}}^{B/C}(D) \cong Z_{\mathbb{P}}(D, \sigma)/N(\sigma)$$

for each $\sigma \in S^{\circ}(\mathbb{Z}_k)$. Take $\sigma = \varepsilon$, the identical permutation. If $\mu \in N(\varepsilon)$, then, according to (4),

$$\mu(i) - i \in \mathbb{Z}_k$$

does not depend on $i = 1, \dots, k$, that is, μ is a power of the cycle

$$(12 \dots k).$$

This fact gives us the required isomorphism

$$\mathbb{P}_{\text{alg}}^{B/C}(D) \cong Z_{\mathbb{P}}(D, \varepsilon)/\mathbb{Z}_k \cong Z_{\mathbb{P}}^{\text{an}}(D)/\mathbb{Z}_k \cong \mathbb{P}_{\text{an}}^{B/C}(D),$$

since the coincidence $Z_{\mathbb{P}}(D, \varepsilon) = Z_{\mathbb{P}}^{\text{an}}(D)$ is obvious. It is easy to check that the isomorphism just constructed is compatible with the isomorphisms

$$\mathbb{P}_{\text{alg}}^{B/C}(D^{\sharp}) \cong \mathbb{P}_{\text{an}}^{B/C}(D^{\sharp})$$

that were constructed above for open subsets $D^{\sharp} \subset D \cap C^{\circ}$.

Q.E.D.

2.5 Divisors of fiber-wise type $(M, \dots, M)$ on $\mathbb{P}^{B/C}$

In this subsection we construct the special model $X^{\sharp}/C$ which was already discussed in sect. 1.3. Recall the rational map

$$p: X \times_C B \dashrightarrow \mathbb{P},$$

regular over C° . Using the projection onto the second factor, we extend p to the map

$$p_B: X \times_C B \dashrightarrow \mathbb{P}_B = \mathbb{P} \times B,$$

also regular over C° . Now construct the product

$$p_B^{\times_C k}: \underbrace{(X \times_C B) \times_C \dots \times_C (X \times_C B)}_k \dashrightarrow \underbrace{\mathbb{P}_B \times_C \dots \times_C \mathbb{P}_B}_k$$

and note that there is a natural isomorphism

$$(X \times_C B)^{\times_C k} \cong X^{\times_C k} \times_C B^{\times_C k}.$$

Taking the composition with the diagonal embedding

$$\Delta: X \hookrightarrow X^{\times_C k},$$

we get the rational map

$$p_B^{\times_C k} \circ (\Delta, \text{id}): X \times_C B^{\times_C k} \dashrightarrow \mathbb{P}_B^{\times_C k}.$$

It is easy to see that this map is equivariant with respect to the action of the symmetric group S_k and that the image of $X \times_C Z_B$ lies in $Z_{\mathbb{P}}$. Factorizing by S_k and taking into consideration that Z_B/S_k is just C , we get a map

$$\iota: X \dashrightarrow \mathbb{P}^{B/C},$$

where $\iota|_{X(D)} = \iota_D: X(D) \rightarrow \mathbb{P}^{B/C}(D)$ for any $D \subset C^\circ$. The image

$$\iota(X) = X^\sharp \subset \mathbb{P}^{B/C}$$

is the desired special model of X/C .

Thus we have realized (on the birational level) the fiber space X/C as a hypersurface $X^\sharp \subset \mathbb{P}^{B/C}$ of fiber-wise type $(M, \dots, M)$. Now let us give an explicit description of such hypersurfaces.

Fix a system of homogeneous coordinates $(u_0, \dots, u_M)$ on $\mathbb{P}$. Let

$$\mathbf{M} = \{\mathcal{M} = u_0^{a_0} u_1^{a_1} \dots u_M^{a_M} \mid a_0 + \dots + a_M = M\} = \{\mathcal{M}_j \mid j \in J\}$$

be the set of all monomials of degree M in the variables u_* . For every point $x \in C^\circ$ the fiber $\tau^{-1}(x) = \{y_1, \dots, y_k\}$ consists of precisely k points, and the fiber $\mathbb{P}_x^{B/C}$ naturally identifies with the product $\mathbb{P}^{(1)} \times \dots \times \mathbb{P}^{(k)}$, where $\mathbb{P}^{(i)} = \mathbb{P}$ corresponds to the point $y_i \in B$. Consider the set of all monomials of the type $(M, \dots, M)$ on the space $\mathbb{P}_x^{B/C}$:

$$\mathbf{M}^{\tau^{-1}(x)} = \{\mathcal{M}_{i_1}^{(1)} \dots \mathcal{M}_{i_k}^{(k)} \mid i_l \in J\},$$

where the monomial $\mathcal{M}_{i_l}^{(l)}$ is in the variables $u_j^{(l)}$, the homogeneous coordinates on the space $\mathbb{P}^{(l)}$. Obviously, there are

$$\sharp(\mathbf{M}^{\tau^{-1}(x)}) = (\sharp \mathbf{M})^k$$

such monomials.

Defining a hypersurface of the type $(M, \dots, M)$ in $\mathbb{P}^{\times k} = \mathbb{P}_x^{B/C}$ is equivalent to give a point in the projective space

$$\mathbb{P}^{(\sharp M)^k - 1}$$

or a non-zero set of coefficients

$$\{c_{i_1, \dots, i_k} \in \mathbb{C} \mid i_l \in J\} / \mathbb{C}^*$$

modulo a non-zero factor. For a given hypersurface of the type $(M, \dots, M)$ we can change the system of coordinates and make sure that for some index $e \in J$

$$c_{e, e, \dots, e} \neq 0.$$

Normalizing this coefficient to 1, we determine all the other coefficients $c_{i_1, \dots, i_k}$.

In particular, for our hypersurface $X^\sharp \subset \mathbb{P}^{B/C}$ we obtain:
at a point $x \in C^\circ$ of general position the set of coefficients, defining the hypersurface $X_x^\sharp$ in $\mathbb{P}_x^{B/C}$, is unique. It is easy to see that the functions on C° that we get as coefficients are branches of multi-valued rational algebraic functions on C . More precisely, let $D \subset C^\circ$ be a simple open set

$$\tau^{-1}(D) = \coprod_{i=1}^k D_i \subset B.$$

Then we get a unique set of meromorphic functions

$$c_{j_1, \dots, j_k} : D \rightarrow \mathbb{C},$$

defining the hypersurface $X(D)$:

$$X(D) = \left\{ \sum_{(j_1, \dots, j_k) \in J^k} c_{j_1, \dots, j_k} \mathcal{M}_{j_1}^{(1)} \dots \mathcal{M}_{j_k}^{(k)} = 0 \right\}.$$

Now we have to globalize the meromorphic functions $c_\sharp$ as branches of multi-valued rational functions on C , that is, rational functions on certain covers of the curve C . Obviously, if

$$c_{i_1, \dots, i_k} \quad \text{and} \quad c_{j_1, \dots, j_k}$$

are two branches of one and the same multi-valued meromorphic function, then the k -uples

$$(i_1, \dots, i_k) \quad \text{and} \quad (j_1, \dots, j_k)$$

differ by a permutation of the symbols only. More formally, for any partition

$$k = k_1 + \dots + k_r$$

and any set

$$\mathcal{M}_1, \dots, \mathcal{M}_r$$

of pair-wise distinct monomials from $\mathbf{M}$ consider all the maps

$$\varepsilon: \tau^{-1}(x) \rightarrow \mathbf{M}, \quad \sharp \varepsilon^{-1}(\mathcal{M}_i) = k_i.$$

Such maps and only such maps (for a fixed partition and a fixed set of pair-wise distinct monomials $\mathcal{M}_i$) can generate different branches of the same multi-valued function.

Let

$$\alpha_{k_1, \dots, k_r}: A_{k_1, \dots, k_r} \rightarrow C$$

be the finite cover, parametrizing for any point $x \in C^\circ$ of general position all the partitions of the fiber $\tau^{-1}(x)$ into a system of subsets of cardinality $k_1, \dots, k_r$:

$$A_{k_1, \dots, k_r} = Z_B / (S_{k_1} \times \dots \times S_{k_r}),$$

where $S_{k_1} \times \dots \times S_{k_r} \hookrightarrow S_k$ is the natural inclusion. The curve $A_{k_1, \dots, k_r}$ certainly can be reducible. In this case when we speak about a rational function

$$a_{k_1, \dots, k_r}: A_{k_1, \dots, k_r} \rightarrow \mathbb{C}$$

we mean an arbitrary set of rational functions on its components.

A hypersurface of fiber-wise type $(M, \dots, M)$ in $\mathbb{P}^{B/C}$ is defined by a set of rational functions

$$a_{k_1, \dots, k_r}^{\mathcal{M}_1, \dots, \mathcal{M}_r}: A_{k_1, \dots, k_r} \rightarrow \mathbb{C}$$

for all the partitions $k = k_1 + \dots + k_r$, $k_1 \geq \dots \geq k_r \geq 1$, and all the r -uples of pair-wise distinct monomials $\mathcal{M}_1, \dots, \mathcal{M}_r$ from $\mathbf{M}$. The sets

$$\left\{ a_{k_1, \dots, k_r}^{\mathcal{M}_1, \dots, \mathcal{M}_r} \right\} \quad \text{and} \quad \left\{ b_{k_1, \dots, k_r}^{\mathcal{M}_1, \dots, \mathcal{M}_r} \right\}$$

define the same hypersurface if and only if there exists a non-zero rational function

$$\lambda: C \rightarrow \mathbb{C}$$

such that

$$a_{k_1, \dots, k_r}^{\mathcal{M}_1, \dots, \mathcal{M}_r} = (\lambda \circ \alpha_{k_1, \dots, k_r}) b_{k_1, \dots, k_r}^{\mathcal{M}_1, \dots, \mathcal{M}_r}$$

for all $(k_1, \dots, k_r)$, $(\mathcal{M}_1, \dots, \mathcal{M}_r)$.

Fixing a set of rational functions $a_\sharp^\sharp$, which defines the hypersurface $X^\sharp = \iota(X) \subset \mathbb{P}^{B/C}$, our description of the special model $X^\sharp \subset \mathbb{P}^{B/C}$ is complete: everything is ready for investigation of the problem of existence of a section of the morphism $\pi: X^\sharp \rightarrow C$.

3 Existence of sections

In accordance with our strategy, we look at the sections of the fibration $X^\sharp/C$ as the sections of the ambient fibration $s: C \rightarrow \mathbb{P}^{B/C}$, $\pi \circ s = \text{id}_C$, the image of which lies in $X^\sharp$, i.e. $s(C) = X^\sharp$.

3.1 Sections of the ambient fibration $\mathbb{P}^{B/C}$

Let $x \in C^\circ$. To give a section $s: C \rightarrow \mathbb{P}^{B/C}$ means to give a value

$$s(x) \in \mathbb{P}_x^{B/C} = \prod_{b \in \tau^{-1}(x)} \mathbb{P}_b$$

for every point $x \in C^\circ$, which is the same as to give values

$$\{s_B(b) \in \mathbb{P}_b | b \in \tau^{-1}(x)\}.$$

Hence sections of the ambient fibration are in one-to-one correspondence with sections

$$s_B: B \rightarrow \mathbb{P}_B = \mathbb{P} \times B.$$

In their turn, the latter sections are determined by $(M+1)$ -uples of rational functions on B .

Now we introduce certain finite-dimensional spaces of sections in the following way. Let R be an effective divisor on B . Set

$$\mathcal{S}(R) = \mathcal{L}(R)^{M+1} = \{(s_0, \dots, s_M) | s_i \in \mathcal{L}(R) = H^0(\mathcal{O}_B(R))\}.$$

Obviously,

$$\dim \mathcal{S}(R) = (M+1) \dim \mathcal{L}(R)$$

and any non-zero $s \in \mathcal{S}(R) \setminus \{0\}$ determines a section of the fibration $\mathbb{P}_B/B$ and thus of $\mathbb{P}^{B/C}/C$. If $\deg R \gg 0$ is, then the map

$$\mathbb{P}(\mathcal{S}(R)) \rightarrow \text{Sect}(\mathbb{P}_B/B) = \text{Sect}(\mathbb{P}^{B/C}/C)$$

is injective on a non-empty Zariski open subset of $\mathbb{P}(\mathcal{S}(R))$.

From now on we assume that $\tau(R) \subset C^\circ$, so that R does not contain ramification points of the morphism τ .

3.2 Sections of the fibration $X^\sharp/C$

Let $x \in C^\circ$, $D \ni x$ be a simple open set, so that

$$\tau^{-1}(D) = \prod_{i=1}^k D_i,$$

$\tau: D_i \rightarrow D$ are isomorphisms for all i . Consider

$$\mathbb{P}^{B/C}(D) = \mathbb{P}^{(1)} \times \dots \times \mathbb{P}^{(k)} \times D,$$

and let $u_j^{(i)}$ be the homogeneous coordinates on $\mathbb{P}^{(i)}$, which come from the coordinates $u_\sharp$ on $\mathbb{P}$. In terms of these coordinates the hypersurface $X^\sharp(D) \subset \mathbb{P}^{B/C}(D)$ is given by the equation

$$F_D(u_i^{(1)}; \dots; u_i^{(k)}; z) = \sum_{(i_1, \dots, i_k) \in J^k} c_{i_1, \dots, i_k}(z) \mathcal{M}_{i_1}^{(1)} \dots \mathcal{M}_{i_k}^{(k)},$$

where the monomials $\mathcal{M}_i^{(j)}$ are in the variables $u_{\sharp}^{(j)}$ and of degree M , whereas the meromorphic functions $c_{\sharp}(z)$ are branches of the functions

$$a_{k_1, \dots, k_r}^{\mathcal{M}_1, \dots, \mathcal{M}_r},$$

with z a coordinate on D . The single-valued branches exist because all the maps

$$\alpha_{k_1, \dots, k_r} : A_{k_1, \dots, k_r} \rightarrow C$$

are reducible over D :

$$\alpha_{k_1, \dots, k_r}^{-1}(D) = \coprod_{\beta \in \mathcal{B}(k_1, \dots, k_r)} D_{\beta},$$

where $\mathcal{B}(k_1, \dots, k_r)$ is the set of all partitions of the type $(k_1, \dots, k_r)$ of the set $\{1, \dots, k\}$, parametrizing the points of the fiber $\tau^{-1}(x)$.

Take $s = (s_0, \dots, s_M) \in \mathcal{S}(R)$. Obviously,

$$s|_{\tau^{-1}(D)} = \{s|_{D_i} = (s_0^{(i)}, \dots, s_M^{(i)}) | i = 1, \dots, k\}.$$

The image of the section $s_D : D \rightarrow \mathbb{P}^{B/C}(D)$, defined by $s \in \mathcal{S}(R) \setminus \{0\}$, lies in $X^{\sharp}(D)$ if and only if

$$F_D(s_{\sharp}^{(1)}; s_{\sharp}^{(2)}; \dots; s_{\sharp}^{(k)}; z) \equiv 0.$$

Let us understand what happens when $s_{\sharp}^{(i)}$ are substituted as $u_{\sharp}^{(i)}$ in the equation F . Let $s \in \mathcal{S}(R)$ be an arbitrary element (possibly zero).

Proposition 3.1 *The meromorphic functions of the variable z*

$$F_D(s_D) = F_D(s_{\sharp}^{(1)}; \dots; s_{\sharp}^{(k)}; z)$$

glue together into a single-valued rational algebraic function $F(s)$ on C .

Proof. This is obvious, since both $s_j^{(i)}(z)$ and $c_{\sharp}(z)$ come from the globally (over C) defined objects: the branches of multi-valued algebraic functions on C , and for $D_1 \subset D_2 \subset C^{\circ}$ we get $F_{D_1}(s_{D_1}) = F_{D_2}(s_{D_2})|_{D_1}$, so that $F_D(s_D)$ glue together into a single-valued function on C° and thus on C . Q.E.D.

3.3 The crucial fact

Let $(F(s))$ be the divisor of poles and zeros of the rational function $F(s)$ on C .

Proposition 3.2 *There exists an effective divisor Δ on the curve C , depending on the hypersurface $X^{\sharp}$ only (but not depending on R), such that for $s \in \mathcal{S}(R)$*

$$(F(s)) + M\tau_*R + \Delta \geq 0. \tag{5}$$

Proof. We observe that up to a divisor Δ which does not depend on R the poles of the rational function $F(s)$ are bounded from above by the divisor

$$M\tau_*R = M\left(\sum_{i \in I} m_i \tau(y_i)\right), \quad \text{where} \quad R = \sum_{i \in I} m_i y_i.$$

Let us prove first a local version of the estimate (5). Let $D \subset C^\circ$ be a simple open set,

$$\tau^{-1}(D) = \coprod_{i=1}^k D_i,$$

$\tau: D_i \rightarrow D$ is an isomorphism, z is a coordinate on D , $z_i = \tau^*z|_{D_i}$ is a coordinate on D_i . Consider the meromorphic function

$$\begin{aligned} F_D(s_D) &= F_D(s_\#^{(1)}; \dots; s_\#^{(k)}; z) = \\ &= \sum_{(i_1, \dots, i_k) \in J^k} c_{i_1, \dots, i_k}(z) \mathcal{M}_{i_1}(s_\#^{(1)}) \dots \mathcal{M}_{i_k}(s_\#^{(k)}), \end{aligned}$$

where $s_j^{(i)}(z) = (\tau^{-1})^*(s_j|_{D_i})$.

The local version of (1) is now provided by

Proposition 3.3 *There exists an effective divisor Δ^+ on C° , depending on $X^\sharp$ only, such that for any simple open set $D \subset C^\circ$*

$$(F_D(s_D)) + M\tau_*(R \cap \tau^{-1}(D)) + \Delta^+ \cap D \geq 0.$$

Proof. Since the degree of the monomials $\mathcal{M}_i$ is equal to M , whereas by construction for any i, j

$$(s_j^{(i)}) + (R \cap D_i) \geq 0,$$

we get for any $j \in J$

$$(\mathcal{M}_j(s_\#^{(i)})) + M(R \cap D_i) \geq 0.$$

This implies

$$\left(\prod_{e=1}^k \mathcal{M}_{i_e}(s_\#^{(e)})\right) + M\tau_*(R \cap \left(\prod_{e=1}^k D_i\right)) \geq 0,$$

that is,

$$\left(\prod_{e=1}^k \mathcal{M}_{i_e}(s_\#^{(e)})\right) + M\tau_*(R \cap \tau^{-1}(D)) \geq 0.$$

Recall that the functions $c_{i_1, \dots, i_k}$ are branches of the rational functions

$$a_{k_1, \dots, k_r}^{\mathcal{M}_1, \dots, \mathcal{M}_r}: A_{k_1, \dots, k_r} \rightarrow \mathbb{C}$$

and thus depend on $X^\sharp$ only. Their poles are fixed and therefore there exists an effective divisor Δ^+ on C° such that

$$(F_D(s_D)) + M\tau_*(R \cap \tau^{-1}(D)) + \Delta^+ \cap D \geq 0,$$

which is what we need.

Q.E.D.

Now we globalize (3.3) and prove Proposition 3.2 From the local fact which we have just obtained we immediately get the estimate

$$(F(s)) \cap C^\circ + M\tau_*R + \Delta^+ \geq 0$$

(since $\text{Supp } \tau_*R \subset C^\circ$ by assumption). Consequently, to complete the proof of Proposition 3.2 it remains to consider the poles of the function $F(s)$ at the points $x \in C \setminus C^\circ$ and prove that they are bounded. But this is easy and done as follows.

In order to be able to work with coordinates, we use the analytic description of $\mathbb{P}^{B/C}$ near a point $x \in C \setminus C^\circ$ given above. Let $D \ni x$ be a small disk, then

$$\mathbb{P}^{B/C}(D) = Z_{\mathbb{P}}(D)/G_1 \times \dots \times G_a.$$

Let us pull back all the functions to $Z_{\mathbb{P}}^{\text{an}}(D)$. The polynomial F_D outside the point x , that is, on $D^\sharp = D \setminus \{x\}$, is defined by the construction, given above. Since by assumption $R \subset C^\circ$, all the functions $s_j^{(i)}$ are regular on $\tau^{-1}(D)$, and therefore the monomials $\mathcal{M}_j$ in these functions are regular, too. The polynomial F_D is linear in its coefficients, which depend on $X^\sharp$ only. Thus there exists an effective divisor $\Delta^\sharp$ with the support in $C \setminus C^\circ$,

$$\Delta^\sharp = \sum_{x \in C \setminus C^\circ} m_x x,$$

such that near $x \in C \setminus C^\circ$ we have

$$(F_D(s)) + (\Delta^\sharp \cap D) \geq 0.$$

Setting $\Delta = \Delta^+ + \Delta^\sharp$, we complete the proof of Proposition 3.2.

Q.E.D.

3.4 End of the proof of Theorem 1.2

As we have seen above, the map

$$\varphi: s \mapsto F(s) \in \mathbb{C}(C)$$

is a map between affine varieties

$$\varphi: \mathcal{S}(R) \rightarrow \mathcal{L}(M\tau_*R + \Delta).$$

By construction (locally over each simple open subset $D \subset C^\circ$ and therefore globally due to the well-known characterization of polynomial functions) φ is polynomial of degree Mk , that is, makes a regular map of affine spaces

$$\varphi: \mathbb{C}^{\dim \mathcal{S}(R)} \rightarrow \mathbb{C}^{\dim \mathcal{L}(M\tau_*R + \Delta)}.$$

Obviously $0 \in \varphi^{-1}(0)$, so that the fiber $\varphi^{-1}(0)$ over the point $0 \in \mathcal{L}(M\tau_*R + \Delta)$ is non-empty. Consequently,

$$\begin{aligned} \dim \varphi^{-1}(0) &\geq \dim \mathcal{S}(R) - \dim \mathcal{L}(M\tau_*R + \Delta) \\ &\quad \parallel \\ &\geq \deg R - K, \end{aligned} \tag{6}$$

where $K \in \mathbb{Z}$ is a constant. Therefore if $\deg R \gg 0$ is sufficiently large, then the fiber $\varphi^{-1}(0)$ contains non-zero elements, which define sections of the fibration $X^\sharp/C$.

The proof of Theorem 1.2 is now complete.

The general theory [KMM96] now implies that the fibration $X^\sharp/C$ has plenty of sections, connecting arbitrary points $w_i \in X^\sharp$ in distinct fibers. The map φ constructed above gives an explicit description of sections as solutions of a system of algebraic equations, whereas the inequality (6) gives an estimate for the dimension of the space of sections with bounded poles. The concluding part of our arguments is a generalization of the classical proof of the Tsen theorem for the “twisted” (not locally trivial) fibration $\mathbb{P}^{B/C} \rightarrow C$.

4 Fibrations of Fano threefolds

Our first result is

Theorem 4.1 *Let $\pi : X \rightarrow C = \mathbb{P}_1$ be a fibrations with general fiber X_b a Fano 3-fold with $b_2(X_b) = 1$. Then π has a section.*

Proof. We are going to apply the classification of Fano 3-folds and refer to [IP99] with the further references given there. Let r be the index of X_b .

If $r = 4$ or $r = 3$, then X_c is $\mathbb{P}_3$ or Q_3 , and the claim follows from Theorem 1.2.

If $r = 2$, then X_b is a del Pezzo 3-fold and all cases are again settled by 1.2 except for X_b being of type V_5 , i.e. X_b is a linear section of $G(2, 5)$ embedded by Plücker in $\mathbb{P}_9$. In that case we consider the relative Douady space $\tau : \mathcal{D} \rightarrow B$ of lines in π -fibers. Since the surface of lines in X_c is just $\mathbb{P}_2$, the map τ is a generic $\mathbb{P}_2$ -fibration over B . Thus τ has a section providing a single rational curve in the general fiber X_c . These rational curves define a rational surface in X over $\mathbb{P}_1$, so that π has a rational section.

It remains to treat the case $r = 1$. Let g be the genus of X_c . If $g \leq 5$, then again X_c is a complete intersection in a weighted projective space, so we can apply 1.2. So let $g \geq 6$; here we will use the Mukai construction. Let $C^* \subset C$ be the set of regular values of π and fix $c \in C$. Then there exists a unique stable vector bundle $\mathcal{E}_c$ over X_c which is spanned and whose sections define a map

$$\phi_c : X_c \rightarrow G(m, V_c)$$

where $V_c = H^0(\mathcal{E}_c)$ such that $\mathcal{E}_c = \phi^*(E)$ for the tautological bundle E on $G(m, V_c)$. Note that ϕ_c is an embedding if $g \geq 7$ but it may be a two-sheeted cover if $g = 6$. Moreover $\phi_b(X_c)$ is a complete intersection in $G(m, V_c)$ if $g \geq 8$. Now every stable bundle on X_c of the same rank and same Chern classes of $\mathcal{E}_c$ is isomorphic to $\mathcal{E}_c$. The $\mathcal{E}_c$ clearly glue locally; and we are going to show that they even glue globally to a vector bundle $\mathcal{E}$ over $X^* = \pi^{-1}(C^*)$. In fact, choose an open covering $U_i \subset C^*$ such that $\mathcal{E}_i$ exists over $V_i = \pi^{-1}(U_i)$. Then the uniqueness of $\mathcal{E}_c$ yields isomorphisms

$$\phi_{ij} : \mathcal{E}_i|_{V_{ij}} \rightarrow \mathcal{E}_j|_{V_{ij}}.$$

By stability, we have $\text{Aut}(\mathcal{E}_c) = \mathbb{C}^*$, hence the cocycle condition

$$\phi_{ij}\phi_{jk} = \phi_{ik}$$

is satisfied up to $\mathbb{C}^*$. This means that at least that the $\mathbb{P}(\mathcal{E}_i)$ glue to a projective bundle over X^* . Since X is a rationally connected fibration over $\mathbb{P}_1$, this bundle has to come from a vector bundle $\mathcal{E}$.

As a consequence we obtain a map

$$\psi : X \rightarrow Z$$

to a fibration $\sigma : Z \rightarrow C$ which is a Grassmann bundle over C^* and which is birationally a product. Moreover $\psi|_{X_c} = \phi_c$ for $c \in C^*$. If now $g = 8$ or $g = 6$ and X is not special, then we can apply the next proposition (Tsen for Grassmann bundles) to conclude. The special case (with genus 6) is the following: the sections of $\mathcal{E}_c$ provide a 2-sheeted covering

$$\phi_c : X_c \rightarrow G(2, 5)$$

the image of which is a Fano 3-fold of type V_5 . Now the (analytic) preimage of a line in V_5 is a conic in X_c and conversely every conic in X_c is mapped to a line in V_5 . Therefore the conics in X_c form an irreducible family, parametrized by a surface S , which admits a map $h : S \rightarrow \mathbb{P}_2$, the surface of line in V_5 . Clearly two smooth conics cannot be mapped to the same line, hence h is birational, i.e. S is rational. Now the same argument as in the case V_5 provides a section in X .

In the remaining cases more efforts are necessary, given in 4.3 and 4.5 below. Q.E.D.

Proposition 4.2 *Let $G = G(p, N)$ be a Grassmannian (projective notation), let B be a smooth rational curve or even smooth curve of any genus g and let $H \subset B \times G$ are relative smooth complete intersection of elements*

$$H_i \in |\mathcal{O}_G(k_i) \otimes g^*(\mathcal{L}_i)|.$$

If $\sum k_i$ is smaller as the index of G , i.e. $\sum k_i < N + 1$, then $H \rightarrow B$ has a section.

Proof. For simplicity of notations we treat only the case where H is a hypersurface and leave the trivial amplification to the general case to the reader.

A section s of $H \rightarrow B$ is given by $p+1$ elements $s_i \in H^0(B, (\mathcal{O}_B(d_i))^{\oplus(N+1)})$, where $0 \leq i \leq N+1$, subject to the following conditions:

1. $s_0(b), \dots, s_p(b)$ are linearly independent for general $b \in B$;
2. $s_0(b) \wedge \dots \wedge s_p(b) \in H_b$ for all $b \in B$.

Set

$$V(d_*) := \bigoplus_{i=0}^p H^0(\mathcal{O}_B(d_i))^{\oplus(N+1)}$$

to be the space parametrizing the $(p+1)$ -uples of elements s_i and consider the map

$$\begin{aligned} \pi : V(d_*) \rightarrow \bigoplus_{\substack{I \subset \{1, \dots, N\} \\ |I| = p+1}} H^0(\mathcal{O}_B(d)) &=: P_d \end{aligned}$$

which is given by Plücker coordinates and where we set

$$d = \sum_{i=0}^p d_i.$$

Its image may well be non-closed. Denote by $W(d_*)$ its closure in P_d . The first problem is to compute its dimension $\dim W(d_*)$. From now on we will assume that $d_0 = d_1 = \dots = d_p = e$, so that $d = (p+1)e$. It is enough to prove our claim for one particular choice of degrees d_i , so let us take this one. In this case denote $V(d_*)$ by V_e , $W(d_*)$ by W_e . Note that the morphism

$$\pi : V_e \rightarrow W_e$$

is dominant by construction.

Lemma. *The relative dimension*

$$\dim V_e/W_e = \dim V_e - \dim W_e = (p+1)^2$$

as soon as $e \geq 2g+1$.

Putting off the proof of the lemma for a while, let us complete the proof of (4.2). Indeed, if the hypersurface H is defined by the equation $F \in H^0(\mathcal{O}_G(k) \otimes \pi^*(\mathcal{L}))$, then the condition (2) is rewritten in Plücker coordinates as

$$F(s(b)) = \sum_{|\alpha|=k} p_\alpha(b) s^\alpha = 0.$$

A section $s = (s_I)$ of the trivial fibration $B \times G/B$ given by its Plücker coordinates s_I is actually a section of the fibration H/B if and only if $F(s(b)) \equiv 0$. Now assume

that $s \in W_d$. so that $s_I \in H^0(\mathcal{O}_B(d))$ for each $I \subset \{1, \dots, N\}$. It is clear that the map

$$s \mapsto F(s) \in H^0(\mathcal{O}_B(kd + m))$$

where $m = \deg(\mathcal{L})$, determines a polynomial map

$$\varphi: W_d \rightarrow H^0(\mathcal{O}_B(kd + m))$$

of closed affine algebraic varieties. Now $\varphi^{-1}(0) \ni 0$, so that the fiber over 0 is non-empty. Thus

$$\begin{aligned} \dim \varphi^{-1}(0) &\geq \dim W_d - h^0(\mathcal{O}_B(m + kd)) \sim \\ &\sim \dim V_d - kd \sim (N + 1 - k)d \end{aligned}$$

up to a bounded integer. Thus, precisely as in the usual Tsen theorem, we obtain a section. Q.E.D.

for (4.2).

Proof of the lemma. We have to show that if $(s_0, \dots, s_p)$ is as above and generic and if $(s'_0, \dots, s'_p)$, satisfies

$$s'_0 \wedge \dots \wedge s'_p = s_0 \wedge \dots \wedge s_p;$$

so that

$$s' = As \tag{+}$$

with a $(p + 1) \times (p + 1)$ -matrix $A = (a_{ij})$ of meromorphic functions on B with $\det A = 1$, then A is constant, i.e. all a_{ij} are constant functions. To this extent we define a meromorphic map

$$\psi: B \times \mathbb{P}_p \rightarrow \mathbb{P}_N$$

by setting

$$\psi(b, [a_0 : a_1 : \dots : a_p]) = [a_0 s_0 + \dots + a_p s_p(b)] = [\sigma(b)].$$

Here the bracket $[\cdot]$ on the right hand side means the projection from a vector space $V \setminus 0$ to projective space, resp. $[\sigma(b)] = [\sigma_0(b) : \dots : \sigma_N(b)]$, where the σ_j are the components of $\sigma = \sum a_i s_i$. Now we will show

Claim (4.2.2) If $e \geq 2g + 1$ is sufficiently large, then ψ is holomorphic and birational onto its image.

First we show how to deduce our lemma from (4.2.2). For simplicity of notation, let $\mathcal{G} = \mathcal{O}_B(e)$. Consider again $s' \in H^0(B, \mathcal{G}^{\oplus N+1})$ such that $s' = As$ with $\det A = 1$. We shall write $s' = \sum a_j s_j$. If $e \geq 2g + 1$, then $\mathcal{G}$ is very ample and therefore the map $\Phi_{s''}: B \rightarrow \mathbb{P}_N$ mapping b to $[s''(b)]$, is an embedding for general $s'' \in H^0(B, \mathcal{G}^{\oplus N+1})$ and the degree of the curve $\Phi_{s''}(B) \subset \mathbb{P}_N$ is independent of s'' . Since s is generic and since we only have to compute the dimension of the fiber at the general point,

we may assume that s' is general in the sense that the previous sentences apply to s' instead of s'' . Since $s' = As$, we conclude that $\Phi_{s'}(B) \subset W$, the image of the birational map ψ . So if B' denotes the image of the map

$$B \rightarrow B \times \mathbb{P}_p, \quad b \mapsto (b, [s'(b)]),$$

then $\Phi_{s'}(B) = \psi(B')$. Now in homology we can write

$$[B'] = [B \times x] + a[l]$$

with a line $l \subset \mathbb{P}_p$ and some integer $a \geq 0$. Let $H = \psi^*(\mathcal{O}_{\mathbb{P}_N}(1))$. Then $H \cdot B' = H \cdot B + a$. On the other hand, $H \cdot B' = \deg \Phi_{s'}(B) = \deg \Phi_s(B) = H \cdot B$. Therefore $a = 0$ and there exists $a = [a'_0 : \dots : a'_p] \in \mathbb{P}_p$ such that $[s'] = [\sum a'_j s_j]$. Hence there is a meromorphic function a on B such that

$$s' = a(\sum a'_j s_j),$$

i.e. $aa'_j = a_j$ for all j . We want to argue that then a must be constant. If not, then s' would always have zeroes. But s' is a general section of a direct sum of very ample line bundle over a curve and therefore has no section.

Proof of (4.2.2) If $d' \geq 2g + 1$, then the evaluation map

$$\epsilon : H^0(\mathcal{G}^{\oplus N+1}) \rightarrow \mathcal{G}_b^{\oplus N+1} \oplus \mathcal{G}_{b'}^{\oplus N+1}$$

is surjective for $b, b' \in B$. This includes as usually the case $b = b'$, in which case the sum is understood as $\mathcal{G}_b^{\oplus N+1} \otimes \mathcal{O}_{B,b}/m_b^2$. Therefore we obtain an embedding

$$\tilde{\psi} : B \times \mathbb{P}_p \rightarrow \mathbb{P}_{N'}$$

with $N' = h^0(B, \mathcal{G}^{\oplus N+1}) - 1$. Recall that - possibly after dualizing the Grassmannian - we may assume that

$$2(p+1) \leq N+1.$$

Now let $\tau : \mathbb{P}_{N'} \rightarrow \mathbb{P}_N$ be the projection from a generic $\mathbb{P}_M$ with $M = N' - N - 1$. Then τ is holomorphic since $N+1 > p+1$. Moreover ψ is of the form $\psi = \tau \circ \tilde{\psi}$ and ψ is birational unless every $\mathbb{P}_{N'-N}$ meets $\tilde{\psi}(B \times \mathbb{P}_p)$ at least twice. This can happen only if

$$p+1 = \dim B \times \mathbb{P}_p \geq \text{codim}_{\mathbb{P}_N} \mathbb{P}_{N'-N},$$

i.e. if $p+1 \geq N$. Since $N \geq 2(p+1) - 1$, as stated above, it follows $p = 0$ and $N = 1$, a trivial case. This finishes the proof of (4.2). Q.E.D.

We are now attacking the cases $g = 9, 10, 12$. The key is (4.3); here it is more convenient to use affine notations for the Grassmannians.

Proposition 4.3 *Let $G = G(p, m)$ be the Grassmannian of p -dimensional linear subspaces in $\mathbb{C}^m$. Let $i : G \rightarrow \mathbb{P}^{\binom{m}{p}-1}$ be the Plücker embedding and B be a smooth rational curve. Let $F \subset B \times \mathbb{P}^{\binom{m}{p}-1}$ be a relative linear subspace. Let $\sigma_1, \dots, \sigma_k$ be relative skew-symmetric l -linear forms on $B \times \mathbb{C}^m$ satisfying simultaneously one of the following conditions*

1. $l = 2$;
2. $l = p - 1$ and $k = 1$.

Let $\Sigma \subset B \times G$ be the locus of p -planes which are σ_j -isotropic for $1 \leq j \leq k$. Let $Z = F \cap \Sigma$; we assume that Z is of the expected dimension and smooth over the general point of B . Furthermore let $n = \dim Z - 1$. Assume finally that in case (1) we have

$$m + n > p(m - p) - \binom{p-1}{2} k$$

and that in case (2) we have

$$m + n > p(m - p) - 1.$$

Then the projection $Z \rightarrow B$ has a section.

(4.4) We first show how to derive Theorem 4.1 in cases $g = 9, 10, 12$ from Proposition 4.3. The first part of the proof of Theorem 4.1 gives a bimeromorphic map $\psi : X \rightarrow Z$ with $Z \subset C \times G \subset C \times \mathbb{P}^{\binom{m}{p}-1}$. Here the relevant data are:

- $(m, p) = (6, 3)$ if $g = 9$;
- $(m, p) = (7, 5)$ if $g = 10$,
- $(m, p) = (7, 3)$ if $g = 12$.

Moreover from Mukai's classification, see [IP99, 5.2.3], Z is of the form $Z = F \cap \Sigma$ for appropriate F and Σ with corresponding data (in the notation of 4.3):

- $(l, k) = (2, 1)$ if $g = 9$,
- $(l, k) = (4, 1)$ if $g = 10$
- $(l, k) = (2, 3)$ if $g = 12$.

Now it is just a matter of direct calculation to verify the inequalities we need to apply 4.3.

Proof of Proposition 4.3 The section s we are looking for will be given by a p -tuple of sections $(s^j)_{1 \leq j \leq p} : C \rightarrow \mathbb{C}^m$, given by relative polynomials s^j on C of degree d - at the end we will take limits as $d \rightarrow \infty$. So our section s depends on $mp(d+1)$ parameters.

Case (1) ($l = 2$): The section s we are looking for is subject to the following conditions:

$$\sigma_i(s^j, s^l) = 0$$

for $1 \leq i \leq k$ and $1 \leq j < l \leq p$. Asymptotically in d these are $k \frac{p(p-1)}{2}$ equations, bilinear in the entries s^j . This leads to $\frac{p(p-1)}{2}$ polynomials of degree $2d$, thus to

$\sim k(p-1)pd$ conditions (the vanishing of the corresponding coefficients). These are the conditions which force a section $s : C \rightarrow C \times G$ to map to Σ . To make sure that s maps to Z we need further conditions. The number of linear conditions describing the relative linear space F is asymptotically equal to

$$\dim \Sigma - \dim F = p(m-p) - k \frac{p(p-1)}{2} - n + 1.$$

Since the Plücker coordinates are polynomials of degree p in the s^j , these linear conditions lead to the vanishing of $p(m-p) - k \frac{p(p-1)}{2} - n + 1$ polynomials in x (coordinate on $\mathbb{P}_1$), each of degree pd , and so we get

$$\sim (p(m-p) - k \frac{p(p-1)}{2} - n + 1)pd$$

conditions.

In total we obtain asymptotically

$$kp(p-1)d + (p(m-p) - k \frac{p(p-1)}{2} - n + 1)pd$$

conditions, while we have mpd parameters to our disposal. Hence we get a section as soon as we have more parameters than conditions, and this leads immediately to inequality (1) in (4.3).

Notice that we should also have checked that $s^1 \wedge \dots \wedge s^p \neq 0$. But this is done in exactly the same way as in (4.2).

Case 2: $l = p-1, k = 1$. Here the arguments are completely analogous. We again have mpd parameters. The number of conditions forcing s to be in Σ is asymptotically $dpl = p(p-1)d$, because the dimension of $(p-1)$ -linear alternating forms on $\mathbb{C}^p$ is $p = \binom{p}{p-1}$, and each coefficient of the relative form is a polynomial of degree $p-1$ in the s^j , which leads to a polynomial of degree $p(p-1)d$ in x .

The number of conditions forcing s to be in F is as in case (1)

$$(\dim \sigma - n)pd = (p(m-p) - p - n + 1)pd,$$

and by the inequality (2) we have at least as many parameters as equations so that s exists. Q.E.D.

(4.5) We are discussing the case $g = 7$. Here $m = 9, p = 4$ whereas $l = 2; k = 1$. In that case the inequalities are not satisfied, so there is no conclusion. However we can still conclude that $\Sigma \rightarrow B$ has at least two sections (actually a 1-dimensional family), say s_1 and s_2 . We then choose a base e_i of $\mathbb{C}^9$ carefully so that the equations of $\Sigma \subset C \times G \subset C \times \mathbb{P}^N$ take a nice form.

Using these two sections, we choose the e_i such that s_1 is the subspace generated by $e_1, \dots, e_4$ and $q(e_i, e_j) = \delta_{ij}$. Moreover we may choose $e_6, \dots, e_9$ such that s_2 is given by the 4-plane defined by those e_i . We can also achieve that $q(e_i, e_{5+j}) = 0$ and

$q(e_{5+i}, e_{5+j}) = \delta_{ij}$. Finally let e_5 be in the orthogonal complement of the e_i already chosen. Denoting by x again the variable on C and denoting by $s_{i,j}$ the Plücker coordinates of the section s ($1 \leq i \leq 4$; $1 \leq j \leq 9$), we can write the equations for a section s to be in Z in the following form:

$$a_i(x)s_{i,5}^2 + b_i(t)s_{i,5+i} = 0$$

for $1 \leq i \leq 4$ and

$$a_{ij}(x)s_{i,5}^2 + c_{ij}(x)s_{i,5+j} + d_{ji}(x)s_{j,5+i}$$

for the obvious set of i, j (these are $\binom{4}{2} = 6$ equations). If we associate to $s_{i,5}$ the weight $\frac{d}{2}$ and to all other coordinates weight d , then we obtain, adding the 7 additional linear equation for describing $Z \subset \Sigma$, in total $10d + 7d = 17d$ conditions. On the other hand we have $16d + 4\frac{d}{2} = 18d$ parameters, which is bigger than the number of conditions. This settles the case $g = 7$.

It remains to treat Fano fibrations whose fibers have $b_2 \geq 2$. This is rather tedious and works by running through the classification of Mori and Mukai. Of course the geometry of the Mori contractions plays the decisive role. Here is just one argument as example - details will be worked out in another paper. Suppose that some (hence every) smooth fiber X_c admits only one birational contraction. Then we can contract fiberwise to obtain $X \rightarrow X' \rightarrow C$. If the birational contraction of X_c contracts the divisor to a point, then X' has a section. Otherwise we get a distinguished curve $C_c \subset X'_c$, i.e. we obtain a ruled surface $S \subset X' \rightarrow C$, which has a section.

References

- [Do80] Dolgachev, I.: Newton polyhedra and factorial rings. J. pure and appl. algebra 18, 253-258 (1980)
- [Do81] Dolgachev, I.: Weighted projective spaces. Lecture Notes in Math. 956, 34-71. Springer 1981
- [GHS01] Graber, T.; Harris, J.; Starr, J.: Families of rationally connected varieties. Preprint, June 2001
- [IP99] Iskovskikh, V.A.; Prokhorov, Yu.G.: Fano varieties. Encyclopaedia in Math. Sci. 47. Springer 1999
- [Ts36] Tsen, C.: Quasi-algebraisch abgeschlossene Funktionenkörper. J. Chinese Math. 1, 81-92 (1936)
- [Ko96] Kollár J.: Rational curves on algebraic varieties. Springer 1996
- [KMM92] Kollár J., Miyaoka, Y. and Mori S.: Rationally connected varieties. J. Alg. Geom. 1, 429-448 (1992)

Frédéric Campana Département de mathématiques Université de Nancy I BP 239, 54506 Vandœuvre les Nancy Cedex France frederic.campana@antares.iecn.u-nancy.fr	Thomas Peternell Mathematisches Institut Universität Bayreuth 95440 Bayreuth Germany thomas.peternell@uni-bayreuth.de
--	--

Aleksandr Pukhlikov Steklov Mathematical Institute Gubkina 8 117966 Moscow Russia pukh@mi.ras.ru	Mathematisches Institut Universität Bayreuth 95440 Bayreuth Germany pukh@btm8x5.mat.uni-bayreuth.de
---	---